

Diagnosis of interatrial shunts and the influence of patent foramen ovale on oxygen desaturation in obstructive sleep apnea

Akademisk avhandling
som för avläggande av medicine doktorsexamen
vid Sahlgrenska akademien, Göteborgs universitet kommer att offentlig
försvaras i centralklinikens aula, Sahlgrenska Universitetssjukhuset/Östra
torsdagen den 22 November 2007 kl 9.00

av
Magnus Johansson
Överläkare i Klinisk Fysiologi

Fakultetsopponent:
Professor Eva Nylander,
Universitetssjukhuset Linköping.

Avhandlingen baseras på följande delarbeten.

- I Availability of percutaneous closure for an adult population with interatrial shunts. Johansson M, Söderberg B, Eriksson P. *Cardiology* 2003; 99: 85-89
- II Sizing of Atrial Septal Defects in Adults. Helgason H, Johansson M, Söderberg B, Eriksson P. *Cardiology* 2005; 104: 1-5
- III The influence of patent foramen ovale on oxygen desaturation in obstructive sleep apnoea. Johansson M. C., Eriksson P, Peker Y, Hedner J, Råstam L, Lindblad U. *European Respiratory Journal* 2007; 29: 149-155
- IV Sensitivity for detection of patent foramen ovale increased with increasing number of contrast injections. A descriptive study with contrast transesophageal echocardiography. Johansson M. C., Helgason H, Dellborg M, Eriksson P. *Journal of the American Society of Echocardiography*. E-published ahead of print, October 2007.



GÖTEBORGS UNIVERSITET

Diagnosis of interatrial shunts and the influence of patent foramen ovale on oxygen desaturation in obstructive sleep apnea

Magnus Johansson

Department of Clinical Physiology,
Department of Emergency and Cardiovascular Medicine,
Sahlgrenska University Hospital /Östra, Institute of Medicine,
the Sahlgrenska Academy at Göteborg University, Göteborg, Sweden.

Abstract

Patent foramen ovale (PFO) is found in 27% of the population and although mostly asymptomatic, PFO has been associated with e.g. cryptogenic stroke and, rarely, also with oxygen desaturation. PFO and atrial septal defects may nowadays be closed percutaneously without open heart surgery. Obstructive sleep apnea is a common condition, characterised by cessation of ventilation due to collapse of the upper airways and oxygen desaturation of varying degree.

The availability for percutaneous closure was studied in 66 consecutive patients with an indication for closure of an interatrial shunt and 58 % of the patients were found to be available.

A descriptive study on 51 consecutive patients with atrial septal defect hypothesised that balloon sizing of the defect during percutaneous closure can be replaced by the size measured with pre-catheterisation transesophageal echocardiography. The results showed that the differences between measurements were too large for substituting pre-catheterisation size for balloon sizing.

The influence of PFO on the frequency of oxygen desaturations in proportion to the frequency of ventilation disturbances in obstructive sleep apnea was studied in a case control study. The presence of a PFO was assessed with contrast transesophageal echocardiography and ≥ 20 bubbles passing over to the left atrium was considered as a large PFO. The prevalence of large PFOs was 9 out of 15 (60%) cases with frequent desaturations, versus only 2 out of 15 controls (13%) ($p=0.02$) with infrequent desaturations, in proportion to the frequency of ventilation disturbances.

The effect of increasing numbers of contrast injections during transesophageal echocardiography, on the sensitivity for PFO detection, was studied. The sensitivity increased with increasing numbers of contrast injections and to safely rule out the presence of a PFO, up to five contrast injections were needed.

In conclusion, interatrial shunts can often be closed percutaneously and balloon sizing is an important part of the procedure. Nocturnal oxygen desaturation occurred proportionally more often in obstructive sleep apnea subjects with a PFO than in subjects without a PFO, indicating the importance of right-to-left shunting in obstructive sleep apnea subjects with a concomitant PFO. Furthermore, sensitivity for PFO detection increased with increasing numbers of contrast injections during transesophageal echocardiography.

Keywords: Patent foramen ovale, atrial septal defect, obstructive sleep apnea, contrast transesophageal echocardiography, percutaneous closure.